



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 35/78, 39/385, 31/35, 31/205, 31/12		A1	(11) International Publication Number: WO 97/02041
			(43) International Publication Date: 23 January 1997 (23.01.97)
(21) International Application Number: PCT/US96/11270		(81) Designated States: AU, BR, CA, JP, MX, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 3 July 1996 (03.07.96)		Published <i>With international search report.</i>	
(30) Priority Data:			
60/000,842 3 July 1995 (03.07.95) US 60/005,643 19 October 1995 (19.10.95) US 08/676,095 2 July 1996 (02.07.96) US			
(71)(72) Applicant and Inventor: CRANDALL, Wilson, T. [US/US]; Route 616, Jolly Hill, Fl. Defiance, VA 24437 (US).			
(74) Agents: STULTS, Larry, W. et al.; Jones & Askew, 37th floor, 191 Peachtree Street, N.E., Atlanta, GA 30303-1769 (US).			
(54) Title: TRANSDERMAL AND ORAL TREATMENT OF ANDROGENIC ALOPECIA			
(57) Abstract			
<p>This invention relates to the topical and oral treatment of hair loss, especially androgenic alopecia, by providing formulations that include anti-androgens, especially extracts of the saw palmetto plant, co-enzyme Q, and acetyl carnitine, and optionally stimulators of adenylate cyclase to stimulate hair growth, to increase the luster of hair, and to decrease hair graying.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic			SE	Sweden
CG	Congo	KR	Republic of Korea	SG	Singapore
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LR	Liberia	SZ	Swaziland
CS	Czechoslovakia	LT	Lithuania	TD	Chad
CZ	Czech Republic	LU	Luxembourg	TG	Togo
DE	Germany	LV	Latvia	TJ	Tajikistan
DK	Denmark	MC	Monaco	TT	Trinidad and Tobago
EE	Estonia	MD	Republic of Moldova	UA	Ukraine
ES	Spain	MG	Madagascar	UG	Uganda
FI	Finland	ML	Mali	US	United States of America
FR	France	MN	Mongolia	UZ	Uzbekistan
GA	Gabon	MR	Mauritania	VN	Viet Nam

TRANSDERMAL AND ORAL TREATMENT OF ANDROGENIC ALOPECIA

Prior Related Applications

This application claims priority to United States provisional Patent Applications Serial Numbers 60/000,842 and 60/005,643 filed on July 3, 1995 and October 19, 1995, respectively.

Technical Field of the Invention

This invention relates to the topical and oral treatment of hair loss, especially androgenic alopecia, by providing formulations that include anti-androgens, especially extracts of the saw palmetto plant, co-enzyme Q, and acetyl carnitine, and optionally stimulators of adenylate cyclase, to stimulate hair growth, to increase the luster of hair, and to decrease hair graying.

Background of the Invention

Androgenic alopecia is an autosomal disorder which begins in puberty in genetically disposed individuals. Androgenic alopecia is also known as hereditary baldness, male pattern baldness, and seborrheic alopecia and occurs in males and females. The disorder is heterogeneous and increased circulating androgens are not the only causative factor. Historically, patients with male androgenic alopecia present with frontal recession of the hairline, especially at the temples and vertex, and androgenetic frontal hairline (incipient

regression of terminal hairs into shorter thin hairs of the intermediate and vellous type). Minoxidil, available since 1988, produces a maximum of only 40% cosmetic responses in selected patients with vertex balding who are young, recently diagnosed and display small areas of alopecia. The response to minoxidil is not seen for 4 to 10 months and treatment must be maintained or the hairline regresses.

Saw palmetto is a small, creeping palm (*Serenoa repens*) of the southeastern United States, having palmately divided leaves with one-ribbed segments and black, one-seeded fruit. It is a native American tree of South Carolina and Georgia and extracts of this tree have been used successfully to treat benign prostatic hypertrophy. Extracts of saw palmetto act as a multi-site inhibitor of the hormone dihydrotestosterone (DHT) which is responsible for prostatic hypertrophy. Saw palmetto blocks approximately 50% of the binding of DHT to receptors in the prostate. It also blocks the uptake of DHT into the nucleus of prostatic cells, and strongly inhibits the action of the enzyme testosterone 5 alpha-reductase which reduces the conversion of testosterone to DHT.

Methods and compositions for increasing the growth of hair are needed, especially in situations of hair loss due to androgenic alopecia. One composition should be easy to apply topically and should promote hair growth. Another composition should be orally administered and promote hair growth.

Summary of the Invention

In one embodiment, the present invention provides an easy to use topical therapeutic composition and treatment for increasing the growth of hair.

In another embodiment, the present invention provides a therapeutic composition and oral treatment for increasing the growth of hair and the luster of hair, and for decreasing the graying of hair.

Accordingly, it is an object of the present invention to provide a treatment for hair loss through the topical application of anti-androgens in various formulations designed for topical application.

5 It is an object of the present invention to provide a treatment for hair loss through the topical application of extracts of saw palmetto, which act as anti-androgens, in various formulations designed for topical application.

10 Another object of the invention to provide a treatment for hair loss through the oral application of anti-androgens in various formulations designed for oral application, combined with co-enzyme Q and acetyl carnitine.

15 It is another object of the invention to provide a treatment for hair loss through the oral application of saw palmetto extracts in various formulations designed for oral application, combined with co-enzyme Q and acetyl carnitine.

20 Another object of the invention to provide a treatment for hair loss through the oral application of saw palmetto extracts in various formulations designed for oral application, combined with co-enzyme Q, acetyl carnitine, and stimulators of adenylate cyclase.

25 Still another object of the invention is to provide a composition and treatment for hair loss through the topical application of anti-androgens combined with co-enzyme Q and acetyl carnitine in various formulations designed for topical application.

30 Another object of the invention is to provide a composition and treatment for hair loss through the topical application of anti-androgens combined with stimulators of adenylate cyclase, co-enzyme Q and acetyl carnitine in various formulations designed for topical application.

Still another object of the invention is to provide a composition and treatment for hair loss through the topical application of extracts of saw palmetto combined with co-

enzyme Q and acetyl carnitine in various formulations designed for topical application.

These and other objects, features and advantages of the present invention will become apparent after a review of the following detailed description of the disclosed embodiments.

Detailed Description of the Invention

The following United States provisional Patent Applications are incorporated by reference herein in their entirety, Serial Numbers 60/000,842 and 60/005,643 filed on July 3, 1995 and October 19, 1995, respectively.

The term "saw palmetto" refers to a native American tree often found in the southeastern part of the United States, especially in Georgia and Florida. As used in the present context, the term "saw palmetto" includes extracts of the saw palmetto tree. These extracts may be present as oil extracts among other forms.

The term "androgen" refers to testosterone and its precursors and metabolites, and 5-alpha reduced androgens, including but not limited to dihydrotestosterone. Androgen refers to androgens from the testis, adrenal gland, and ovaries, as well as all forms of natural, synthetic and substituted or modified androgens.

It is to be understood that other agents which affect the synthesis, metabolism, site of action, and function of androgens are considered within the scope of the present invention when combined with acetyl carnitine and co-enzyme Q. Included within the scope of this invention are drugs, extracts, chemicals, or other agents which affect the biosynthesis, enzymatic conversion, binding to receptors, and metabolism of androgens. Agents which modify the translocation of androgens to the nucleus or the second messenger systems that transduce intracellular signals are also

considered within the scope of this invention when combined with acetyl carnitine and co-enzyme Q. Agents which affect the synthesis, release, and function of hypothalamic and pituitary hormones which affect the synthesis or release of testosterone are also included in the scope of this invention when combined with acetyl carnitine and co-enzyme Q for stimulating hair growth. Accordingly, methods for stimulating hair growth which employ acetyl carnitine and co-enzyme Q combined with agents described in this paragraph which affect the synthesis, release, efficacy, or metabolism of androgens are considered within the scope of this invention. Accordingly, molecules which affect hypothalamic synthesis and secretion of gonadotropin releasing hormone (GnRH), GnRH agonists (such as leuprolide and gonadorelin) which affect pituitary release of gonadotropins, ketoconazole and liarazole which affect testosterone synthesis in the testis, finasteride and saw palmetto extract which affect conversion of testosterone to dihydrotestosterone by 5-alpha reductase in its various forms in extraglandular tissues, and flutamide and cyproterone acetate which affect binding of androgen to its receptor, are all considered within the scope of this invention. The aforementioned molecules in this paragraph are collectively defined as "anti-androgens" within this application.

The term "androgenic alopecia" refers to an autosomal disorder which begins in puberty in genetically disposed individuals. Androgenic alopecia is also known as hereditary baldness, male pattern baldness, and seborrheic alopecia. Androgenic alopecia may occur in males and females.

The term "acetyl carnitine" refers to various forms of acetyl carnitine, including but not limited to, D,L-carnitine, and acetyl-L-carnitine hydrochloride which is usually employed in a 1% to 5% solution. Acetyl carnitine (PCAA, Houston, TX) is thought to decrease in senescence and down regulation. Although not wanting to be bound by this

statement, it is believed that the decrease in acetyl carnitine compromises the transport of fatty acids from the cytosol into the inner mitochondrial membrane of the mitochondria, thereby impairing the physiological function of the follicle, perhaps through effects on fatty acid oxidation and ATP production. Acetyl carnitine is believed to improve fatty acid metabolism by stimulation of cardiolipin which affects the activity of the inner mitochondrial membrane, affecting its permeability and function for proton transport. Acetyl carnitine is also believed to synergize with co-enzyme Q in the revitalization of senescent hair follicles.

The term "co-enzyme Q" refers to several forms of co-enzyme Q including co-enzyme Q0, co-enzyme Q2, co-enzyme Q6, co-enzyme Q7, co-enzyme Q9 and co-enzyme Q10, and are all considered within the scope of this invention. Several of these forms are also known as ubiquinones and may be obtained from PCAA (Houston, TX). Various forms of co-enzyme Q (PCAA, Houston, TX) are utilized because it is believed that the senescent hair follicles are metabolically reduced through reduced activity of respiratory enzymes. Although not wanting to be bound by this statement, it is believed that co-enzyme Q, especially co-enzyme Q10, increases respiratory levels compatible with hair formation. Although not wishing to be bound by this statement, it is believed that supplementation with co-enzyme Q facilitates removal of excess protons accumulated at the inner mitochondrial membrane due to prolonged down regulation, acts as an antioxidant, and promotes normalization of oxidative function (Shigenaga, M. *et al.* Proc. Soc. Natl. Acad. Sci. USA 99: 10771-10778, 1994).

The term "lecithin" is used to mean a phosphatide called phosphatidylcholine and can be isolated from soybean, eggs and other sources including, but not limited to, heart, brain, and liver. Lecithin is a mixture of the diglycerides of stearic, palmitic, and oleic acids, linked to the choline ester of

phosphoric acid. Soybean lecithin is a preferred lecithin and may contain the following acids; palmitic, stearic, palmitoleic, oleic, linoleic, linolenic and arachidonic as described in the *Merck Index* (Ninth edition, 1976, page 711).

5 The term "PLURONIC" refers to poloxamer compounds and are sold collectively under the trademark PLURONIC (BASF, Parsippany, NJ). PLURONIC F-127 corresponds to poloxamer 407, a polyoxypropylene-polyoxyethylene block copolymer described by Schmolka in the *Journal of Biomedical Materials Research* 6: 571-582, 10 (1972). As used in this application, the terms PLURONIC organogel, poloxamer organogel, and polyoxyethylene/polyoxypropylene organogel are synonymous.

 "Topical" application is used to mean local 15 administration of the composition and its various embodiments, for example, in the treatment of alopecia. The composition according to the present invention can be in the form of solutions, lotions, salves, creams, ointments, liposomes, sprays, gels, roller sticks or any other method using 20 micelles and pharmaceutically acceptable penetration enhancers. In one embodiment, the composition may be applied to the scalp at bedtime and again after showering in the morning for a total of two applications per day.

 The "enhanced penetration" caused by 25 compositions of this invention as used in topical application with this method, means increased penetration into the skin, and is achieved with compounds such as lecithin organogel, PLURONIC organogel (BASF, Parsippany, NJ), N-decylmethyl sulfoxide (NDMS), and ethoxydiglycol:ethanol, and can be 30 observed in many ways known to those skilled in the art.

 "Oral administration" refers to the administration 35 of the compositions of the present invention into the gastrointestinal tract in any form, including but not limited to the following, tablets, powders, suspensions, solutions, lozenges, and lollipops.

The present invention includes a composition for topical treatment of androgenic alopecia comprising anti-androgens, including but not limited to saw palmetto extracts, combined with co-enzyme Q and acetyl carnitine. These compounds are delivered topically combined with delivery vehicles and penetrating agents optionally containing lecithin, isopropyl palmitate, lecithin organogel, PLURONIC F-127 organogel, NDMS, ethoxydiglycol:ethanol and/or water. When co-enzyme Q is used in the absence of organogels, it is first dissolved in ethoxydiglycol:ethanol and then in Tween, such as Tween 20 or Tween 80, glycerol, castor oil, olive oil, or another appropriate agent. For example, these agents are useful in the formation of the composition of the present invention as a roller for topical application.

In one embodiment, therapeutically effective amounts of the following compounds may be combined at the indicated concentrations expressed per 120 ml of formulation for topical application: saw palmetto (1 g to 20 g), co-enzyme Q (0.1 g to 10 g), and acetyl carnitine (0.1 g to 20 g). Preferred concentrations of these compounds for use in topical application are provided in Example 4.

Another preferred penetrating agent and delivery vehicle is lecithin organogel which is a combination of lecithin, isopropyl palmitate and water. Lecithin organogels have been described as vehicles that are useful in facilitating the delivery of low molecular weight compounds transdermally (Willmann, H., et al., "Lecithin Organogel as Matrix for Transdermal Transport of Drugs", *J. Pharm. Sci.*, Vol. 81, 1992, which is incorporated herein by reference). The lecithin organogels are obtained by adding small amounts of water to a solution of lecithin in organic solvents. Generally, lecithin organogels are prepared at room temperature by first dissolving lecithin in an organic solvent such as isopropyl palmitate and then adding enough water while stirring to obtain the desired gel. Preparation of a variety of lecithin

gels, all of which are appropriate in practicing the present invention, are described in Scartazzini, et al. *Journal of Physical Chemistry*, Vol. 92, pgs. 829-833 (1988) and Luisi, P.L. et al. *Colloid and Polymer Science*, Vol. 268, pgs. 356-374 (1990), both of which are incorporated herein by reference. The lecithin organogel preferably comprises 1:1 to 1.5:5 (weight/vol) of soy lecithin (PCAA, Kinghurst, Houston, TX) to isopropyl palmitate (PCAA) (1g:1ml). Other penetrating agents may be used in the composition of the present invention.

The formation of organogels containing lecithin dissolved with isopropyl palmitate or other solvents, and water, has been described by Luisi et al., *Colloid and Polymer Science* 268: 356-374 (1990) and Scartazzini et al., *The Journal of Physical Chemistry* 92: 829-833 (1988). Willimann et al., *Journal of Pharmaceutical Sciences* 81:871-874 (1992), examined the efficacy of lecithin organogels for use in the transdermal delivery of drugs such as scopolamine and broxaterol. Willimann et al., also observed that lecithin organogels had no detrimental effect on skin when compared to control samples treated with physiological saline (see page 872, column 2, paragraph 3, *Journal of Pharmaceutical Sciences* 81:871-874 (1992)).

The present invention optionally includes lecithin (20% - 98%) dissolved with isopropyl palmitate or other solvents in combination with an approximately 10% - 20% solution of PLURONIC F-127 (BASF, Parsippany, NJ), otherwise known as poloxamer 407, in a ratio of approximately 1:3 to 1:4. This ratio may be varied by one of ordinary skill in the art. Other PLURONICS may be used in the present invention. It is to be understood that the soy lecithin of the present invention is a preferred lecithin source and may be dissolved in isopropyl palmitate to achieve a final concentration in the composition of from approximately 20% - 98%, with a more preferred final concentration of from

approximately 20% - 40%. Lecithins may optionally be derived from eggs, and organs such as heart, brain, and liver, and used at concentrations of approximately 20% - 99%, with more preferred final concentrations of from approximately 20% - 40%. The composition according to the present invention can be in the form of lotions, salves, creams, ointments, liposomes, sprays, micelles, or gels. The desired form is lotions, ointments and salves. Liposomes are described in detail by Oleniacz in U.S. Patent No. 3,957,971, the entirety of which is hereby incorporated by reference.

Another delivery vehicle is *n*-Decylmethyl sulfoxide (NDMS obtained from PCAA 10925). NDMS is optionally present at a concentration of between approximately 0.01% and 1% by weight, with a preferred concentration of between approximately 0.05% and 0.5% by weight, with the most preferred concentration of approximately 0.125% by weight. NDMS is dissolved in 10 ml of a 75% solution of ethanol. Next, acetyl carnitine is dissolved in purified water. Co-enzyme Q is added to 2 to 5 ml of ethoxydiglycol and then to 10 ml of 70% ethanol plus Tween 80. Finally, purified water is added to a final volume of 120 ml.

A gelling agent optionally may be added to the formulation. Gelling agents that are suitable for use in the present invention include, but are not limited to, cellulose, cellulose ethers, carboxymethylcellulose, alginates, polyacrylates, bentonite, gelatin, tragacanth, polyvinylpyrrolidone, polyvinyl alcohol, and polyoxyethylene/polyoxypropylene block copolymers, some of which are known as poloxamers. The poloxamer compounds are sold collectively under the trademark PLURONIC (BASF, Parsippany, NJ). PLURONIC F-127 corresponds to poloxamer 407. Other PLURONICS may be used in the present invention. As used in this application, the terms PLURONIC organogel, poloxamer organogel and polyoxyethylene/polyoxypropylene organogel are synonymous.

5 A preservative, such as benzyl alcohol (1%) or potassium sorbate, may be added to the composition. An antioxidant, including but not limited to antioxidants such as vitamin E, proanthocyanidin, or lipoic acid, may be added to lecithin organogels and PLURONIC organogels. Other preservatives well known to those of ordinary skill in the art can be used in the composition.

10 Agents for improving the aroma of the formulation for topical application can optionally be added to the composition. A desired aroma improving agent is honey almond oil (PCAA). Other aroma improving agents include, but are not limited to, avocado oil, sesame oil, castor oil, olive oil, grapeseed oil, clove oil, groundnut oil, corn oil, lemon oil, coconut oil, lime oil, hazelnut oil, jojoba oil, carthamus oil and wheatgerm oil. The oils can be added individually or in combination. It is to be understood that various fragrances and assorted floral scents may be optionally added to the composition and are commercially available (PCAA). Stabilizers, antioxidants, preservatives, humectants, regreasing agents, solvents or auxiliaries can be added to improve stability and/or adhesiveness of the formulations. Cosmetic agents such as panthenol may also optionally be added to the formulation.

25 In addition, antimicrobial agents can be optionally added to the composition of the present invention. Addition of an antimicrobial agent is desirable when treating inflammatory conditions associated with loss of hair.

30 The composition of the present invention can be administered topically twice daily or several times per day depending upon the nature and severity of the condition being treated. The lecithin organogel or PLURONIC organogel may be wiped off about 1 hour after application if desired since the active ingredients are absorbed rapidly.

35 The present invention also includes formulations for the treatment of androgenic alopecia that may be

administered orally. These formulations for oral administration do not contain penetration enhancers such as NDMS or lecithin organogel. The present invention includes a composition for oral treatment of androgenic alopecia comprising anti-androgens such as saw palmetto extracts combined with co-enzyme Q and acetyl carnitine. Typical concentration ranges for these compounds used in oral administration are the following; saw palmetto (10 mg - 1 g), co-enzyme Q (10 mg - 250 mg), and acetyl carnitine (250 mg - 2 g). A preferred composition for administration as a pill into the oral cavity twice daily, at approximately 12 hour intervals, is saw palmetto (160 mg), co-enzyme Q10 (30 mg - 180 mg), and acetyl carnitine (1 g). These compounds for oral administration are delivered into the gastrointestinal tract in any form, including but not limited to the following; tablets, powders, suspensions, solutions, lozenges, lollipops, through catheters, and various tubes, including but not limited to nasogastric, gastric, duodenal, jejunal, ileal and colonic tubes. When administered into the oral cavity, various compounds may be added to improve the flavor and consistency of the composition. For example, the composition could be administered into the oral cavity in the form of a cherry flavored lozenge, or a grape flavored syrup.

While not wanting to be bound by this statement, it is believed that DHT affects adenylate cyclase in the hair follicle, thereby modulating keratin production and affecting hair quality and formation. Accordingly, it is within the scope of this invention to include stimulators of adenylate cyclase in conjunction with anti-androgens to stimulate activity of the hair follicle and increase hair growth. In one embodiment, the drug forskolin, a stimulator of adenylate cyclase, is combined with anti-androgens, such as saw palmetto extracts, co-enzyme Q and acetyl carnitine, and a penetrant for topical application. For topical application, a concentration of about 0.1% to 10% may be used with a preferred concentration of 1% to 5%

forskolin per 120 ml of formulation. In oral preparations, forskolin is included at a concentration of about 0.2 mg to 100 mg, with a preferred concentration of about 2 mg to 20 mg, together with anti-androgens, such as saw palmetto extracts, co-enzyme Q and acetyl carnitine. It is understood that these dosages of forskolin are preferred dosages and that other dosages may suggest themselves to those skilled in the art. Similarly, other stimulators of adenylate cyclase will be administered at dose ranges specific for each stimulator.

This invention is further illustrated by the following examples, which are not to be construed in any way as imposing limitations upon the scope thereof. On the contrary, it is to be clearly understood that resort may be had to various other embodiments, modifications, and equivalents thereof which, after reading the description herein, may suggest themselves to those skilled in the art without departing from the spirit of the present invention.

Example 1

Topical Application of Saw Palmetto Extracts Increase Hair Growth in Males with Alopecia

A 52 year old male at the first recognition of alopecia at the temples and frontal recession of the hairline applied the formulation of the present invention as described below. The composition consisting of the formulation of saw palmetto, co-enzyme Q and acetyl carnitine was used in PLURONIC organogel and NDMS twice a day at the right (R) triangulus for a period of 3 weeks. One morning, large numbers of vellous hairs were evident. Treatment was then extended to the left (L) triangulus where vellous hair growth also occurred. The frontal scalp line was treated with the formulation for a period of 3-4 weeks and intermediate hairs were observed among the anagen hairs. The hair was decisively more mature in appearance. Analysis of root hairs of the frontal scalp line indicated an marked increase in the

percent of telogen follicles to 35% from the normal 20% and an increase in the percent of catogen follicles.

Five individual volunteers from 52 to 56 years of age and another three individual volunteers from 31 to 38 years of age were treated with the topical application of saw palmetto extracts in the composition described in the preceding paragraph. Alopecia in these individuals varied from a rating of Hamilton 2 to Hamilton 8. Totally bald scalp devoid of pigmented hair from 5 to 25 years responded to the disclosed treatment of this invention in an interval of time directly correlated with age, rapidity of hair loss and duration of baldness. In general terms, younger men who were bald for less time responded to the treatment faster than individuals such as older men who were bald for longer periods. Photographs of the scalp of these individuals verified increased hair growth.

Example 2

Lecithin Organogel Composition Prepared With Pluronic

A preferred composition for delivery of the formulation of saw palmetto, co-enzyme Q and acetyl carnitine was prepared as described below. The lecithin organogel was prepared by dissolving 20 g of soy lecithin granules (PCAA) in 20 ml of isopropyl palmitate (PCAA). The mixture was stirred periodically for 24 hours until the soy lecithin granules were dissolved. The PLURONIC gel 20% stock solution was prepared by dissolving 16 g of PLURONIC F127 powder (BASF, Parsippany, NJ), also called poloxamer 407, in 80 ml of purified water. Potassium sorbate (160 mg; PCAA) was added to the PLURONIC gel 20% stock solution as a preservative. This was placed in the refrigerator at about 4°C for about 24 hours and stirred periodically.

The composition was prepared by mixing 20 ml of the lecithin organogel with 2 ml of the honey almond oil (PCAA) until a smooth mixture was prepared. Honey almond

oil was added for fragrance. Next, 80 ml of the PLURONIC gel 20% stock solution was mixed in until a gel formed. A blender was used for this mixing step at room temperature with disinfected equipment. The gel was stored at room temperature.

Example 3

Lecithin Organogel Composition Prepared Without Pluronic

In another embodiment of the present invention for delivery of the formulation, the composition described in Example 2 was prepared using lecithin organogel without the addition of the PLURONIC gel 20% stock solution. The final concentration of lecithin organogel was in the range of 20 - 45 % by modifying the ratio of lecithin organogel to water.

Example 4

Compositions For The Topical Treatment Of Androgenic Alopecia

Extracts of saw palmetto are added at a concentration of 5 g to 7.5 g per 120 ml of formulation to the compositions of Examples 2 and 3 in effective therapeutic concentrations. Next, therapeutically effective amounts of co-enzyme Q10 (0.5 g to 1 g per 120 ml of formulation) and acetyl carnitine (1 g to 2 g per 120 ml of formulation) are added to this mixture of saw palmetto in the compositions of Examples 2 and 3 to create the formulation for topical treatment of alopecia.

Example 5

Compositions And Methods For The Treatment Of Androgenic Alopecia By Oral Administration

A composition for treatment of androgenic alopecia by oral administration comprises saw palmetto extracts (160 mg), combined with the following components selected from the group comprising co-enzyme Q10 (30 mg to 180 mg) and acetyl carnitine (1 gm).

These compounds are usually delivered twice daily into the gastrointestinal tract in forms including, but not limited to the following; tablets, powders, suspensions, solutions, lozenges, and lollipops, and also through catheters, and various tubes, including but not limited to nasogastric, gastric, duodenal, jejeunal, ileal and colonic tubes. These compounds are delivered into the oral cavity in the form of tablets which may be chewed, dissolved or swallowed or as powders, suspensions, optionally combined with various excipients, flavor enhancers or other compounds to enhance the texture of the composition.

It should be understood, of course, that the foregoing relates only to preferred embodiments of the present invention and that numerous modifications or alterations may be made therein without departing from the spirit and the scope of the invention.

Claims

5 1. A composition for increasing hair growth comprising an anti-androgen, co-enzyme Q, and acetyl carnitine.

10 2. The composition of Claim 1, further comprising a stimulator of adenylate cyclase.

15 3. The composition of Claim 2, further comprising a penetrant, the penetrant comprising ethoxydiglycol and ethanol, isopropyl palmitate, lecithin organogel, soybean organogel, poloxamer organogel, water, or N-decylmethyl sulfoxide.

20 4. The composition of Claim 2, further comprising an effective amount of a preservative, an antimicrobial, a gelling agent, and an aroma improving agent.

25 5. The composition of Claim 2, wherein the anti-androgen is saw palmetto extract.

30 6. The composition of Claim 2, wherein the stimulator of adenylate cyclase is forskolin.

35 7. A method for increasing hair growth comprising administration of an amount of an anti-androgen, co-enzyme Q, and acetyl carnitine, effective to increase hair growth.

8. The method of Claim 7, further comprising
a stimulator of adenylate cyclase.

5

9. The method of Claim 7, wherein the anti-
androgen is saw palmetto extract.

10

10. The method of Claim 9, wherein the
administration is into the gastrointestinal tract and the effective
amount of saw palmetto extract is about 10 mg to 1 g, the
effective amount of co-enzyme Q is about 10 mg to 250 mg,
and the effective amount of acetyl carnitine is about 250 mg to
2 g.

15

11. The method of Claim 10, wherein the
effective amount of saw palmetto extract is about 50 mg to 500
mg, the effective amount of co-enzyme Q is about 30 mg to
180 mg, and the effective amount of acetyl carnitine is about
500 mg to 1.25 g.

20

12. The method of Claim 8, wherein the
administration is into the gastrointestinal tract, wherein the
anti-androgen is saw palmetto extract, the stimulator of
adenylate cyclase is forskolin, and the effective amount of saw
palmetto extract is about 10 mg to 1 g, the effective amount of
co-enzyme Q is about 10 mg to 250 mg, the effective amount
of acetyl carnitine is about 250 mg to 2 g, and the effective
amount of forskolin is about 0.2 mg to 100 mg.

25

30

35

13. The method of Claim 10, comprising administration in tablets, powders, suspensions, solutions, lozenges, lollipops, through catheters, and through tubes, comprising nasogastric, gastric, duodenal, jejunal, ileal and colonic tubes.

14. The method of Claim 9, wherein the administration is topical and the saw palmetto extract, co-enzyme Q and acetyl carnitine are combined with a penetrant in a formulation.

15. The method of Claim 14, further comprising a stimulator of adenylate cyclase in the formulation.

16. The method of Claim 15, wherein the penetrant is ethoxydiglycol and ethanol, isopropyl palmitate, lecithin organogel, soybean organogel, poloxamer organogel, water, or N-decylmethyl sulfoxide.

17. The method of Claim 16, wherein the penetrant is a combination of lecithin organogel and poloxamer organogel in a ratio of about 1 to 4.

18. The method of Claim 14, wherein the amount of saw palmetto extract is about 1 g to 20 g, the amount of co-enzyme Q is about 0.1 g to 10 g, and the amount of acetyl carnitine is about 0.1 g to 20 g per 120 ml of formulation.

19. The method of Claim 14, wherein the amount of saw palmetto extract is about 4 g to 10 g, the amount of co-enzyme Q is about 0.5 g to 2 g, and the amount of acetyl carnitine is about 0.8 g to 4 g per 120 ml of formulation.

20. The method of Claim 15, wherein the stimulator of adenylate cyclase is forskolin, and wherein the amount of saw palmetto extract is about 1 g to 20 g, the amount of co-enzyme Q is about 0.1 g to 10 g, the amount of acetyl carnitine is about 0.1 g to 20 g, and the amount of forskolin is about 0.1% to 10% per 120 ml of formulation.

21. The method of Claim 15, comprising administration in solution, lotion, salve, ointment, cream, micelle, liposome, spray, gel, roller stick, and spray.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/11270**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) : A61K 35/78, 39/385, 31/35, 31/205, 31/12

US CL : 424/195.1; 514/455, 556, 690

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/195.1; 514/455, 556, 690

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Biosis, Medline, Hcaplus, Wpids, Embase- the four types of active agents herein in hair or alopecia treating or dermatological compositions

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,314,689 A (SCANDURRA ET AL.) 24 May 1994, column 4.	1-21
Y	US 4,713,397 A (HIRAMA ET AL.) 15 December 1987, the abstract.	1-21
Y	Database HCAPLUS, American Chemical Society, AN 1991:234867, LAL, B. ET AL. "Pharmaceutical compositions comprising labdane diterpenoid derivatives and pyrimido (6,1-a) isoquinolin-4-one derivatives and their use," abstract, EP 370379 A1, May 1990.	2-21
Y	Database WPIDS, Derwent Information Ltd., AN 90-307329 [41], FR 2643375 A, (BELLANGER, F. ET AL.), abstract.	1-21



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

A

document member of the same patent family

Date of the actual completion of the international search

01 OCTOBER 1996

Date of mailing of the international search report

10 OCT 1996

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks

Box PCT

Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

M. MOEZE

Telephone No.

(703) 308-1235

